



Gene Comprehensive Nutrigenomic Report

Accession Number: #####

Specimen Collected: ###/###/####

Specimen Received: ###/###/####

Report Generated: November 30, 2022


Specimen Type: Buccal Swab

Provider: #####

Patient Name: #####

Patient DOB: ###/###/####

Patient Gender: Male



Do not make any decisions about your health solely based on the information contained in this report.
Always consult with a licensed and experienced health practitioner when you receive this report.

– 34 – Male

(-/-) No clinical abnormality

(+/-) Heterozygous result

(+/+) Homozygous result

rsID	Gene	Genetic Result	Therapeutics Associated With Positive Result	Highly Recommended Therapeutics	Provider Discretion: As Needed Formula Recommendations	Lifestyle Recommendations	Laboratory Recommendations
Methylation / Foundation Panel							
Methylation and Folate Metabolism							
rs2071010	FOLR1	-/-	Methyltetrahydrofolate (5-MTHF)	Methyl Folate Plus™ Twice Daily			
rs651933	FOLR2	+/+					
rs1643649	DHFR	+/-					
rs6495446	MTHFS	+/-					
rs1076991	MTHFD1	+/-					
rs1801131	MTHFR A1298C	+/+					
rs1801133	MTHFR C677T	-/-					
rs1051266	SLC19A1	+/+					
Methylation and B12 Metabolism							
rs1805087	MTR	-/-	L-5-Methyl THF, Methyl Cobalamin, Niacinamide (B3), Methionine	Methylation Pro Topical™ OR Methylation Complete Fast Dissolves™ twice daily			Consider Plasma B12
rs1802059	MTRR A664A	+/-	Methyl B12				
rs1801394	MTRR A66G	+/-					
rs558660	GIF	+/-	Methyl B12, Adenosyl B12				
rs526934	TCN1	+/-					
rs1801198	TCN2	+/+					

– 34 – Male

(-/-) No clinical abnormality

(+/-) Heterozygous result

(+/+) Homozygous result

rsID	Gene	Genetic Result	Therapeutics Associated With Positive Result	Highly Recommended Therapeutics	Provider Discretion: As Needed Formula Recommendations	Lifestyle Recommendations	Laboratory Recommendations
Methylation / Foundation Panel							
Vitamin D Transport							
rs731236	VDR Taq	+/+	High Dose Vitamin D	Vitamin D3+K2 Cofactor Complex™			Consider Vit D
rs2282679	GC or DBP	-/-	Vitamin K				
Mitochondrial							
rs4147730	NDUFS3	-/-	CoQ 10, PQQ, L-Carnitine, Ornithine, Magnesium, NADH, Calcium	Mito Cell PQQ™			
rs809359	NDUFS7	-/-					
rs1051806	NDUFS8	-/-					
rs4850	UQCRC2	+/-					
rs11648723	UQCRC2	-/-					
rs8042694	COX5A	-/-					
rs4626565	COX6C	-/-					
rs1244414	ATP5C1	-/-					
rs6535454	CoQ2	+/+					

Summary for Foundation / Methylation / Wellness

Highly Recommended Therapeutics

- Methyl Folate Plus™ Twice Daily
- Methylation Pro Topical™ OR Methylation Complete Fast Dissolves™ twice daily
- Vitamin D3+K2 Cofactor Complex™
- Mito Cell PQQ™

Provider Discretion: As Needed Formula Recommendations

Lifestyle Recommendations


Laboratory Recommendations

- Consider Plasma B12
- Consider Vit D


Cell

Chromosome

Nucleosome



METHYLATION



Methylation

- Involves the addition of a methyl group (CH₃)
- Regulates gene expression and repression
- Reduces or removes toxins that eliminate essential nutrients
- Provides nutrients needed for processes such as detoxification, immune regulation, gut health

Methionine

- Used in protein formation and stabilization
- Elevated levels are associated with risk for coronary heart disease, stroke & neurological diseases

Glutathione

- Important for chemical detoxification & proper mitochondrial functioning
- Genes relevant for production include: AHCY, CTH, CGTP1, GSTM1, GSTM3, GSR, MTRR & MTR

5-Methyl Folate

- Important for dopamine and serotonin formation, detoxification and mitochondrial strength
- Genes relevant for production include: DHFR, FOLR1/2, MTHFD1, MTHFR, MTHFS

Homocystein

- Elevated levels are associated with risk for coronary heart disease, stroke, neurological diseases
- Variants in the methylation pathway can be associated with increased/decreased levels

FOLATE

FOOD SOURCES



Eggs



Citrus Fruits



Leafy Greens



Legumes



Broccoli




Brussel Sprouts



Breads, Cereal,
Pastas, Rice



**ABSORBABLE
FOLATE**
NATURAL B9



**5-METHYL
FOLATE**
ACTIVE FORM

FUNCTIONS (OR BENEFITS AS YOU AGE)

- Maintains structure & function of proteins
- Maintains structure & function of DNA
- Facilitates DNA replication, neurotransmitter production & detoxification

DEFICIENCY CAUSES

- Neural tube defects
- Cardiovascular disease
- Memory problems
- Depression
- Insomnia
- Irritability

VITAMIN B12

FOOD SOURCES



Eggs



Seafood – clams, trout, salmon, tuna



Meats – liver, beef, ham, chicken



Low-fat milk products



Swiss cheese



Low-fat yogurt



Nori/Seaweed



Fortified Foods



Nutritional Yeast



**ABSORBABLE
VITAMIN B12**



**METHYLCO-
BALAMIN**
ACTIVE FORM

FUNCTIONS (OR BENEFITS AS YOU AGE)

- Formation and maintenance of red blood cells (RBCs)
- Facilitates DNA synthesis
- Regulates homocysteine levels (decreases)
- Facilitates neurological functioning

DEFICIENCY & PATHWAY ALTERATIONS

- Increased production of homocysteine
- Decreased breakdown of homocysteine
- Circadian Rhythm Problems
- Cancers
- Memory-related disorders
- Cardiovascular diseases
- Fatigue
- Poor balance
- Mood disorders

VITAMIN D

FOOD SOURCES



Tuna



Mushrooms



Eggs



Mackerel



Cheese



Milk Products



BENEFITS AS YOU AGE



Lower Risk
of Fractures



Improves
Heart Function



Supports
Immune System



Speeds
Wound Healing

DEFICIENCY CAUSES

- Bone Pain
- Arthritis
- Obesity
- Backache
- Depression
- Diabetes
- Hypertension
- Osteoporosis
- Heart Disease
- Skin Conditions

MITOCHONDRIA

WAYS TO INCREASE MITOCHONDRIAL FUNCTIONING



Build muscle mass
(strength conditioning)



Decrease toxin exposure
(metals, persistent organic pollutants)



Vitamins
Glutathione
CoQ10
α-lipoic acid +
acetyl-L-carnitine
Resveratrol
NAC
Vitamin E
PQQ
Ginkgo biloba
Proanthocyanidins



FUNCTIONS (OR BENEFITS AS YOU AGE)

- Energy production for growth, movement & homeostasis
- Programmed cell death for dysfunctional, old cells
- Calcium signaling for neuron excitability, neurotransmission & plasticity (strengthening)

DEFICIENCY CAUSES

- Early Aging
- Neurological Disorders
- Diabetes
- Developmental Issues
- Psychiatric Disorders
- Cardiovascular Diseases
- Fatigue

Gene Information Key

rsID	Gene	"_" variant	"+" variant
rs1244414	ATP5C1	C	T
rs6535454	CoQ2	A	G
rs8042694	COX5A	A	G
rs4626565	COX6C	T	C
rs1643649	DHFR	T	C
rs2071010	FOLR1	G	A
rs651933	FOLR2	A	G
rs2282679	GC or DBP	T	G
rs558660	GIF	G	A
rs1076991	MTHFD1	C	T
rs1801131	MTHFR:A1298C	T	G
rs1801133	MTHFR:C677T	G	A
rs6495446	MTHFS	C	T
rs1805087	MTR	A	G
rs1802059	MTRR:A664A	G	A
rs1801394	MTRR:A66G	A	G
rs4147730	NDUFS3	G	A
rs809359	NDUFS7	A	G
rs1051806	NDUFS8	C	T
rs1051266	SLC19A1	T	C
rs526934	TCN1	A	G
rs1801198	TCN2	C	G
rs11648723	UQCRC2	G	T
rs4850	UQCRC2	G	A
rs731236	VDR Taq	A	G

Definitions

ESSENTIAL VITAMINS	The polymorphisms in this panel will identify any potential weakness of absorption, conversion or delivery of your essential vitamins.
COQ2	CoQ2 (Para-hydroxybenzoate—polyprenyltransferase, mitochondrial) codes for an enzyme that functions in the final steps in the biosynthesis of CoQ10 (ubiquinone).. This enzyme, which is part of the coenzyme Q10 pathway, catalyzes the prenylation of parahydroxybenzoate with an all-trans polyprenyl group. Mutations in this gene cause coenzyme Q10 deficiency. Polymorphisms in this gene can lead to severe fatigue, muscle weakness, exercise intolerance and general mitochondrial weakness.
GC or DBP	GC aka DBP (Vit. D Binding Protein) gene codes for Vit. D binding protein. This protein belongs to the albumin family and is a multifunctional protein found in plasma, ascitic fluid, cerebrospinal fluid and on the surface of many cell types. It is manufactured in the hepatic parenchymal cells. DBP is capable of binding to all forms of Vit D including ergocalciferol (vitamin D2) and cholecalciferol (vitamin D3), the 25-hydroxylated forms (calcifediol) and the active hormonal product, 1,25-dihydroxyvitamin D (calcitriol). The major proportion of vitamin D in blood is bound to this protein. It transports vitamin D metabolites between skin, liver and kidney, and then on to the various target tissues. It binds to vitamin D and its plasma metabolites and transports them to target tissues. Polymorphisms in this gene decrease the affinity of the protein to Vit. D which reduces the response rate to Vit. D therapy. Patients with these polymorphisms require high doses of Vit D supplementation.
INFLAMMATORY	This Enzyme category has significant effects on the inflammatory state of a person's body. Polymorphisms in these specific enzymes will significantly increase the levels of inflammation in the body. By supplementing these enzyme deficiencies, the patient will effectively reduce inflammatory damage to the body.
VDR Taq1	The Vitamin D (calcitriol) Receptor is a member of the nuclear receptor family. Upon activation by vitamin D (a secosteroid), the VDR causes the activation or deactivation of protein production by the cell. Impaired vitamin D function can result in significant immune weakness and increased cancer risk, as well as, early bone loss, an increased risk of cognitive decline and mood disorders.
METHYLATION	Methylation is a primary biochemical process in the body that involves the addition of a "methyl" chemical group to a vitamin or neurotransmitter. The addition of the "methyl" group allows for very specific biochemical interactions. Poor "methylation" function alters the effectiveness, delivery and function of many vitamins and important chemicals in the cell.
DHFR	Dihydrofolate reductase, or DHFR, is an enzyme that reduces dihydrofolic acid to tetrahydrofolic acid. This enzyme is the second enzyme in the folic acid conversion chain. Having a mutation in this enzyme can create a methylation deficiency with a MTHFR mutation.
FOLR1	Folate Receptor 1 (FOLR1) is a member of the folate receptor (FOLR) family. Members of this gene family have a high affinity for folate. Polymorphisms in this gene allow for poor delivery of folate to the interior of cells. This can create a high plasma folic acid. This polymorphism does create a methylation deficiency. This polymorphism is associated with many disorders of pregnancy.
FOLR2	Folate Receptor 2 (FOLR2) is a member of the folate receptor (FOLR) family. Members of this gene family have a high affinity for folic acid. Polymorphisms in this gene allow for poor delivery of folic acid to the interior of cells. This can create a high plasma folic acid. This polymorphism does create a methylation deficiency. This polymorphism is associated with many disorders of pregnancy. This receptor is found in high quantities on the placenta, thymus and bone marrow. Can be affiliated with immune disorders.
GIF	The glycoprotein product of the Gastric Intrinsic Factor (GIF) gene is secreted by the stomach lining. GIF protein is required for absorption of Vitamin B12. B12 is necessary for normal red blood cell maturation.
MTHFD1	Methylenetetrahydrofolate Dehydrogenase 1 enzyme handles 2 significant enzymes conversions in the production of L-MTHF. This common polymorphism causes a significant methylation deficiency due to the fact that it is utilized in two steps in methyl-folate production.
MTHFR A1298C	Methylenetetrahydrofolate reductase (MTHFR) catalyzes the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, the bioactive form of folic acid. Two significant polymorphism variants exist in this gene, the A1298C and the C677T. The 1298 confers a conversion weakness of 10% for one copy and approximately 20% for two copies. In contrast, the 677 variant is much more severe and conveys a 40% conversion weakness for one copy and 70% for two copies. A reduced level of MTHFolate produces significant biochemical effects including poor production of dopamine and serotonin, pregnancy complications, poor healing of the nervous system, weak mitochondrial function, reduced production of glutathione, poor cell turnover and poor function of T cell lymphocytes.
MTHFR C677T	Methylene tetrahydrofolate reductase (MTHFR) catalyzes the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, the bioactive form of folic acid. Two significant polymorphism variants exist in this gene, the A1298C and the C677T. The 1298 confers a conversion weakness of 10% for one copy and approximately 20% for two copies. In contrast, the 677 variant is much more severe and conveys a 40% conversion weakness for one copy and 70% for two copies. A reduced level of MTHFolate produces significant biochemical effects including poor production of dopamine and serotonin, pregnancy complications, poor healing of the nervous system, weak mitochondrial function, reduced production of glutathione, poor cell turnover and poor function of T cell lymphocytes.
MTHFS	MTHFS (methylenetetrahydrofolate synthase) is an enzyme that catalyzes the conversion of 5-formyltetrahydrofolate to 5,10-methylenetetrahydrofolate, a precursor of reduced folates. This polymorphism codes for a decreased function of the enzyme and results in poor utilization of Leucovorin (5-formyltetrahydrofolate)..

MTR	MTR (Methionine Synthase) codes for the enzyme that catalyzes the final step in methionine biosynthesis. Polymorphisms in this gene lead to poor recycling of methionine from homocysteine. This enzyme work in coordination with MTRR and requires both MTHF and B12 for proper functioning. Deficiencies in Methionine leads to poor methylation that is associated with numerous neurological, cardiovascular and immunological disease states, as well as, infertility and birth defects.
MTRR Ala637=	Methionine Synthase Reductase is an enzyme responsible for the production of methionine, a very important amino acid. Polymorphisms in this enzyme require an increased amount of Methyl B12 to help this reaction.
SLC19A1	The SLC19A1 gene encodes the reduced folate carrier (RFC) protein. Mutations in the RFC are associated with reduced plasma folate.
TCN1	The protein product of the transcobalamin 1 (TCN1) gene binds Vitamin B12 and protects it from the low pH environment of the human stomach. Individuals homozygous for the G allele of the TCN1 SNP, rs526934, are predicted to have lower serum B12.
TCN2	The protein product of the Transcobalamin 2 gene, TCN2, binds the active form of vitamin B-12. Individuals with the G/G phenotype at rs1801198 have decreased serum B-12 and increased homocysteine when compared to individuals with the C/C phenotype.
MITOCHONDRIA	The mitochondrial enzymes are responsible for energy production from the mitochondria. The mitochondria is known as the "powerhouse" of the cell and produces over 90% of the energy for a cell. The mitochondrial respiratory chain (also known as the electron transport chain) is where these 4 protein complexes are found. Polymorphic alterations in these enzymes reduce the energy output of the mitochondria and leads to symptoms of chronic fatigue, cognitive deficiency, exercise intolerance, low metabolic rate, muscle weakness, poor healing and higher rates of sleep disorders and mood abnormalities.
ATP5C1	ATPase 5c1 (ATP5C1) is an enzyme responsible for producing ATP (the energy component) in the mitochondria. This protein is known as Complex V (the 5th protein) in the mitochondrial respiratory chain. Polymorphisms in the gene confer a weakened energy production by the mitochondria.
COX5A	Cytochrome c oxidase subunit 5a (COX5A) is a protein in a subunit of the cytochrome c oxidase complex, also known as Complex IV of the mitochondrial electron transport chain. Polymorphisms in this enzyme produce a weakened energy production by the mitochondria.
COX6C	Cytochrome c oxidase subunit 6c (COX6C) is a protein in a subunit of the cytochrome c oxidase complex, also known as Complex IV of the mitochondrial electron transport chain. Polymorphisms in this enzyme produce a weakened energy production by the mitochondria.
NDUFS3	The NDUFS3 genes encodes a mitochondrial enzyme, NADH Dehydrogenase (Ubiquinone) Fe-S Protein 3. Like other NDUFS proteins, NDUFS3 is thought to require ubiquinone for full activity.
NDUFS7	NADH Dehydrogenase [ubiquinone] iron-sulfur protein 7 (NDUFS7) is a mitochondrial protein also know as Complex I of the mitochondrial respiratory chain. It is located in the mitochondrial inner membrane and is the largest of the five complexes of the electron transport chain. Polymorphisms in this enzyme produce a weakened energy production in the mitochondria.
NDUFS8	NADH Dehydrogenase (Ubiquinone) Fe-S Protein 8 (NDUFS8) encodes an enzyme in the mitochondrial respiratory chain. Mutations in the NDUFS8 gene are associated with Leigh Syndrome, osteoporosis, and mitochondrial complex I deficiency.
UQCRC2 Arg183Gln	Ubiquinol Cytochrome c Reductase (UQCR, Complex II) is a mitochondrial enzyme protein also known as Complex III of the electron transport chain. Polymorphisms in this enzyme produce a weakened energy production by the mitochondria.

Disclaimers

METHODOLOGY AND LIMITATIONS:

Testing for genetic variation/mutation on listed genes was performed using ProFlex PCR and Real-Time PCR with TaqMan® allele-specific probes on the QuantStudio 12K Flex. All genetic testing is performed by GX Sciences, 4150 Freidrich Lane, Ste H, Austin, TX. 78744. This test will not detect all the known alleles that result in altered or inactive tested genes. This test does not account for all individual variations in the individual tested. Test results do not rule out the possibility that this individual could be a carrier of other mutations/variations not detected by this gene mutation/variation panel. Rare mutations surrounding these alleles may also affect our detection of genetic variations. Thus, the interpretation is given as a probability. Therefore, this genetic information shall be interpreted in conjunction with other clinical findings and familial history for the administration of specific nutrients. Patients should receive appropriate genetic counseling to explain the implications of these test results. Details of assay performance and algorithms leading to clinical recommendations are available upon request. The analytical and performance characteristics of this laboratory developed test (LDT) were determined by GX Sciences' laboratory pursuant to Clinical Laboratory Improvement Amendments (CLIA) requirements.

CLIA #: 45D2144988 Laboratory Director: James Jacobson, PhD

DISCLAIMER:

This test was developed and its performance characteristics determined by GX Sciences. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA and qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. rsIDs for the alleles being tested were obtained from the dbSNP database (Build 142).

DISCLAIMER:

UND Result: If you have received the result Variant undetermined (UND) this indicates that we were not able to determine your carrier status based on your raw data. Please refer to the GX Sciences genetic knowledge database for more information: https://www.gxsciences.com/kb_results.asp

DISCLAIMER:

Report contents and report recommendations are created and approved by GX Sciences. Sole responsibility for the proper use of the information on the GX Sciences report rests with the user, or those professionals with whom the user may consult. Nutrigenomic Testing and Dietary Supplements are not "Designated Health Services" covered by Medicare or Medicaid and may not be reimbursed under any state or Federal health care program.

DISCLAIMER:

These products are not approved by the Food and Drug Administration and are not intended to diagnose, treat, cure or prevent disease. These recommendations are for report purposes only and an individual is not required to use such products. These are recommendations only and do not replace the advisement of your own healthcare practitioner.

GX Sciences SNP References

ESSENTIAL VITAMINS SNP References

CoQ2

• Turunen M., Olsson J.; Dallner G.. Metabolism and function of coenzyme Q. *Biochem. Biophys. Acta.* , 2004, vol. 1660 (pg. 171-199) • Santos-Ocana C., Do T.Q., Padilla S., Navas P., Clarke C.F.. Uptake of exogenous coenzyme Q and trnsport to mitochondria is required for bc1 complex sustainability in yeast coq mutants, *J. Biol. Chem.*, 2002, vol. 277 (pg. 10973-10981) • Oh, J., Ban, M. R., Miskie, B. A., Pollex, R. L., & Hegele, R. A. (2007). Genetic determinants of statin intolerance. *Lipids in Health and Disease*, 6. <https://doi.org/10.1186/1476-511X-6-7> • López-Martín, J. M., Salvati, L., Trevisson, E., Montini, G., DiMauro, S., Quinzii, C., ... Navas, P. (2007). Missense mutation of the COQ2 gene causes defects of bioenergetics and de novo pyrimidine synthesis. *Human Molecular Genetics*, 16(9), 1091–1097. <https://doi.org/10.1093/hmg/ddm058> • Jakobs, B. S., Van Den Heuvel, L. P., Smeets, R. J. P., De Vries, M. C., Hien, S., Schaible, T., Rodenburg, R. J. T. (2013). A novel mutation in COQ2 leading to fatal infantile multisystem disease. *Journal of the Neurological Sciences*, 326(1–2), 24–28. <https://doi.org/10.1016/j.jns.2013.01.004> • Hubacek, J. A., Adamkova, V., Zlatohlavek, L., Steiner-Mrazova, L., & Vrablik, M. (2017). COQ2 polymorphisms are not associated with increased risk of statin-induced myalgia/myopathy in the Czech population. *Drug Metabolism and Personalized Therapy*, 32(4), 177–182. <https://doi.org/10.1515/dmpt-2017-0027> • Diomedì-Camassei, F., Di Giandomenico, S., Santorelli, F. M., Caridi, G., Piemonte, F., Montini, G., Emma, F. (2007). COQ2 Nephropathy: A Newly Described Inherited Mitochondriopathy with Primary Renal Involvement. *Journal of the American Society of Nephrology*, 18(10), 2773–2780. <https://doi.org/10.1016/j.jcir.2009.06.001> • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: [https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.](https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple%20carbohydrates%20like%20white%20flour,efficient, long-lasting way.) • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. *Nature Reviews Immunology* (2011). doi:10.1038/nri2975 • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. *Nature* (2014). doi:10.1038/nature12985 • Kann, O. & Kovács, R. Mitochondria and neuronal activity. *American Journal of Physiology - Cell Physiology* (2007). doi:10.1152/ajpcell.00222.2006 • Pizzorno, J. Mitochondria-fundamental to life and health. *Integrative Medicine (Boulder)* (2014).

GC or DBP

• Slater, N. A., Rager, M. L., Havrdá, D. E. & Harralson, A. F. Genetic Variation in CYP2R1 and GC Genes Associated with Vitamin D Deficiency Status. *Journal of Pharmacy Practice* 30, 31–36 (2017). • Hill, A. How to Get Vitamin D: 7 Effective Ways. *Healthline* (2019). Available at: <https://www.healthline.com/nutrition/how-to-increase-vitamin-d> • Database, G. C. H. G. GC Gene (Protein Coding). *GeneCards* Available at: <https://www.genecards.org/cgi-bin/carddisp.pl?gene=GC> • Cheung, C. L., Lau, K. S., Sham, P. C., Tan, K. C. & Kung, A. W. Genetic variant in vitamin D binding protein is associated with serum 25-hydroxyvitamin D and vitamin D insufficiency in southern Chinese. *Journal of Human Genetics* 58, 749–751 (2013). • Davies, J. R., Field, S., Randerson-Moor, J., Harland, M., Kumar, R., Anic, G. M., ... Newton-Bishop, J. (2014). An inherited variant in the gene coding for vitamin D-binding protein and survival from cutaneous melanoma: A BioGenoMEL study. *Pigment Cell and Melanoma Research*. <https://doi.org/10.1111/pcmr.12193> • Elkum, N., Alkayal, F., Noronha, F., Ali, M. M., Melhem, M., Al-Arouj, M., ... Abubaker, J. (2014). Vitamin D insufficiency in Arabs and South Asians positively associates with polymorphisms in GC and CYP2R1 genes. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0113102> • Leong, A., Rehman, W., Dastani, Z., Greenwood, C., Timpson, N., Langsetmo, L., ... Richards, J. B. (2014). The Causal Effect of Vitamin D Binding Protein (DBP) Levels on Calcemic and Cardiometabolic Diseases: A Mendelian Randomization Study. *PLoS Medicine*. <https://doi.org/10.1371/journal.pmed.1001751> • Nimitphong, Hataikarn, Chanika Sritara, La Or Chailurkit, Suwannee Chanprasertyothin, Wipa Ratanachaiwong, Piyamit Sitara, and Boonsong Ongphiphadhanakul. 2015. "Relationship of Vitamin D Status and Bone Mass According to Vitamin D-Binding Protein Genotypes." *Nutrition Journal* 14 (1). BioMed Central Ltd. doi:10.1186/s12937-015-0016-1. • Szkandera, J. et al. Association of common gene variants in vitamin D modulating genes and colon cancer recurrence. *Journal of Cancer Research and Clinical Oncology* 139, 1457–1464 (2013). • Theodoratou, E., Palmer, T., Zgaga, L., Farrington, S. M., McKeigue, P., Din, F. V. N., ... Campbell, H. (2012). Instrumental variable estimation of the causal effect of plasma 25-hydroxy-vitamin D on colorectal cancer risk: A Mendelian randomization analysis. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0037662> • Trummer, O., Langsenlehner, U., Krenn-Pilkso, S., Pieber, T. R., Obermayer-Pietsch, B., Gerger, A., ... Langsenlehner, T. (2016). Vitamin D and prostate cancer prognosis: a Mendelian randomization study. *World Journal of Urology*. <https://doi.org/10.1007/s00345-015-1646-9> • Wang, W., Ingles, S. A., Torres-Mejia, G., Stern, M. C., Stanczyk, F. Z., Schwartz, G. G., ... John, E. M. (2014). Genetic variants and non-genetic factors predict circulating vitamin D levels in Hispanic and non-Hispanic White women: The breast cancer health disparities study. *International Journal of Molecular Epidemiology and Genetics*. • Wang, Y., Wang, O., Li, W., Ma, L., Ping, F., Chen, L., & Nie, M. (2015). Variants in Vitamin D binding protein gene are associated with gestational diabetes mellitus. *Medicine (United States)*. <https://doi.org/10.1097/MD.0000000000001693>

INFLAMMATORY SNP References

VDRTaq

• Adorini, L., Daniel, K. & Penna, G. Vitamin D Receptor Agonists, Cancer and the Immune System: An Intricate Relationship. *Curr. Top. Med. Chem.* (2006). doi:10.2174/156802606777864890 • McKenna, M. J. & Murray, B. Vitamin D deficiency. in *Endocrinology and Diabetes: A Problem-Oriented Approach* (2014). doi:10.1007/978-1-4614-9884-9_23 • Fischer, K. Vitamin D, in *Principles of Nutrigenetics and Nutrigenomics: Fundamentals of Individualized Nutrition* (2019). doi:10.1016/B978-0-12-804572-5.00032-X • Bikle, D. D. Vitamin D metabolism, mechanism of action, and clinical applications. *Chemistry and Biology* (2014). doi:10.1016/j.chembiol.2013.12.016 • Cieslińska, A. et al. Vitamin D receptor gene polymorphisms associated with childhood autism. *Brain Sci.* (2017). doi:10.3390/brainsci7090115 • Kamel, M. M., Fouad, S. A., Salaheldin, O., El-Razek, A. E. R. A. & El-Fatah, A. I. A. Impact of vitamin D receptor gene polymorphisms in pathogenesis of Type-1 diabetes mellitus. *Int. J. Clin. Exp. Med.* (2014). • El-Shal, A. S., Shalaby, S. M., Aly, N. M., Rashad, N. M., & Abdelaziz, A. M. Genetic variation in the vitamin D receptor gene and vitamin D serum levels in Egyptian women with polycystic ovary syndrome. *Mol. Biol. Rep.* (2013). doi:10.1007/s11033-013-2716-y • Keen, R. W., Hart, D. J., Lanchbury, J. S. & Spector, T. D. Association of early, osteoarthritis of the knee with a Taq I polymorphism of the vitamin D receptor gene. *Arthritis Rheum.* (1997). doi:10.1002/art.1780400812 • Kyung, S. P., Jung, H. N. & Choi, J. The short vitamin D receptor is associated with increased risk for generalized aggressive periodontitis. *J. Clin. Periodontol.* (2006). doi:10.1111/j.1600-051X.2006.00944.x • Wang, Y. et al. Variants in Vitamin D binding protein gene are associated with gestational diabetes mellitus. *Med. (United States)* (2015). doi:10.1097/MD.0000000000001693 • Hill, A. How to Get Vitamin D: 7 Effective Ways. *Healthline* (2019). Available at: <https://www.healthline.com/nutrition/how-to-increase-vitamin-d> • Baudino, T. A. et al. Isolation and characterization of a novel coactivator protein, NCoA-62, involved in vitamin D-mediated transcription. *J. Biol. Chem.* (1998). doi:10.1074/jbc.273.26.16434 • Lisse, T. S., Chun, R. F., Rieger, S., Adams, J. S. & Hewison, M. Vitamin D activation of functionally distinct regulatory mRNAs in primary human osteoblasts. *J. Bone Miner. Res.* (2013). doi:10.1002/jbmr.1882 • Fleet, J. C. & Schoch, R. D. Molecular mechanisms for regulation of intestinal calcium and phosphate absorption by vitamin D. in *Vitamin D* (2011). doi:10.1016/B978-0-12-381978-9.10019-8 • Germain, P., Staels, B., Dacquet, C., Spedding, M. & Laudet, V. Overview of Nomenclature of Nuclear Receptors. *Pharmacol. Rev.* (2006). doi:10.1124/pr.58.4.2 • Tagami, T., Lutz, W. H., Kumar, R. & Jameson, J. L. The interaction of the vitamin D receptor with nuclear receptor corepressors and coactivators. *Biochem. Biophys. Res. Commun.* (1998). doi:10.1006/bbrc.1998.9799 • Herdick, M., Steinmeyer, A. & Carlberg, C. Antagonistic action of a 25-carboxylic ester analogue of 1??,25- dihydroxyvitamin D3 is mediated by a lack of ligand-induced vitamin D receptor interaction with coactivators. *J. Biol. Chem.* (2000). doi:10.1074/jbc.M910000199 • Luderer, H. F. & Demay, M. B. The vitamin D receptor, the skin and stem cells. *J. Steroid Biochem. Mol. Biol.* (2010). doi:10.1016/j.jsmb.2010.01.015 • Vdr vitamin D (1,25-dihydroxyvitamin D3) receptor [Mus musculus (house mouse)] - Gene - NCBI. National Center for Biotechnology Information (2018). Available at: <https://www.ncbi.nlm.nih.gov/gene/22337>.

METHYLATION SNP References

DHFR

• Beierlein, J. M., Karri, N. G. & Anderson, A. C. Targeted mutations of bacillus anthracis dihydrofolate reductase condense complex structure-activity relationships. *J. Med. Chem.* (2010). doi:10.1021/jm100727t • Osborne, M. J., Schnell, J., Benkovic, S. J., Dyson, H. J. & Wright, P. E. Backbone dynamics in dihydrofolate reductase complexes: Role of loop flexibility in the catalytic mechanism. *Biochemistry* (2001). doi:10.1021/bi010621k • Chen, M. J. et al. The functional human dihydrofolate reductase gene. *J. Biol. Chem.* (1984). • Funanage, V. L., Myoda, T. T., Moses, P. a & Cowell, H. R. Assignment of the human dihydrofolate reductase gene to the q11---q22 region of chromosome 5. *Mol. Cell. Biol.* (1984). doi:10.1128/mcb.4.10.2010 • Schnell, J. R., Dyson, H. J. & Wright, P. E. Structure, Dynamics, and Catalytic Function of Dihydrofolate Reductase. *Annu. Rev. Biophys. Biomol. Struct.* (2004). doi:10.1146/annurev.biophys.33.110502.133613 • Crabtree, M. J., Tatham, A. L., Hale, A. B., Alp, N. J. & Channon, K. M. Critical role for tetrahydrobiopterin recycling by dihydrofolate reductase in regulation of endothelial nitric-oxide synthase coupling: Relative importance of the de novo biopterin synthesis versus salvage pathways. *J. Biol. Chem.* (2009). doi:10.1074/jbc.M109.041483 • Benkovic, S. J., Fierke, C. A. & Naylor, A. M. Insights into enzyme function from studies on mutants of dihydrofolate reductase. *Science* (80-). (1988). doi:10.1126/science.3125607 • Huennekens, F. M. In search of dihydrofolate reductase. *Protein Sci.* (1996). doi:10.1002/pro.5560050626 • Bailey, S. W. & Ayling, J. E. The extremely slow and variable activity of dihydrofolate reductase in human liver and its implications for high folic acid intake. *Proc. Natl. Acad. Sci.* (2009). doi:10.1073/pnas.0902072106 • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits> • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998> • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Abu Seman, N., Wan Mohamad, W. N., Östenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674

FOR1

• Suzuki, Y., Yoshimoto-Nakagawa, K., Maruyama, K., Suyama, A. & Sugano, S. Construction and characterization of a full length-enriched and a 5'-end-enriched cDNA library. *Gene* (1997). doi:10.1016/S0378-1119(97)00411-3 • Sadasivan E, Rothenberg SP (1989). "Molecular cloning of the complementary DNA for a human folate binding protein" • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Abu Seman, N., Wan Mohamad, W. N., Östenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998> • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>

<https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>. • Henderson, G. B. Folate-binding proteins. *Annu. Rev. Nutr.* (1990). doi:10.1146/annurev.nutr.10.1.319 • Kelemen, L. E. The role of folate receptor ?? in cancer development, progression and treatment: Cause, consequence or innocent bystander? *International Journal of Cancer* (2006). doi:10.1002/ijc.21712 • Ragoussis, J., Senger, G., Trowsdale, J. & Campbell, I. G. Genomic organization of the human folate receptor genes on chromosome 11q13. *Genomics* (1992). doi:10.1016/S0888-7543(05)80236-8 • Elwood, P. C. Molecular cloning and characterization of the human folate-binding protein cDNA from placenta and malignant tissue culture (KB) cells. *J. Biol. Chem.* (1989). • Yan, W. & Ratnam, M. Preferred Sites of Glycosylphosphatidylinositol Modification in Folate Receptors and Constraints in the Primary Structure of the Hydrophobic Portion of the Signal. *Biochemistry* (1995). doi:10.1021/bi00044a039

FOLR2

• Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • Freisheim, J. H., Price, E. M. & Ratnam, M. Folate coenzyme and antifolate transport proteins in normal and neoplastic cells. *Adv. Enzyme Regul.* (1989). doi:10.1016/0065-2571(89)90091-5 • Ragoussis, J., Senger, G., Trowsdale, J. & Campbell, I. G. Genomic organization of the human folate receptor genes on chromosome 11q13. *Genomics* (1992). doi:10.1016/S0888-7543(05)80236-8 • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674 • Nakashima-Matsushita, N. et al. Selective expression of folate receptor beta and its possible role in methotrexate transport in synovial macrophages from patients with rheumatoid arthritis. *Arthritis Rheum.* (1999). doi:10.1002/1529-0131(199908)42:83.0.CO;2-L • Suzuki, Y., Yoshimoto-Nakagawa, K., Mariyama, K., Suyama, A. & Sugano, S. Construction and characterization of a full length-enriched and a 5'-end-enriched cDNA library. *Gene* (1997). doi:10.1016/S0378-1119(97)00411-3 • Page, S. T., Owen, W. C., Price, K. & Elwood, P. C. Expression of the Human Placental Folate Receptor Transcript is Regulated in Human Tissues. *Journal of Molecular Biology* 229, 1175–1183 (1993). • Shen, F., Ross, J. F., Wang, X. & Ratnam, M. Identification of a novel folate receptor, a truncated receptor, and receptor type beta in hematopoietic cells: cDNA cloning, expression, immunoreactivity, and tissue specificity. *Biochemistry* (1994). doi:10.1021/bi00171a021 • Ratnam, M., Marquardt, H., Duhning, J. L. & Freisheim, J. H. Homologous membrane folate binding proteins in human placenta: cloning and sequence of a cDNA. *Biochemistry* (1989). • 5-Methyltetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998>. • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>. • Strausberg, R. L. et al. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc. Natl. Acad. Sci. U. S. A.* (2002). doi:10.1073/pnas.242603899 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Henderson, G. B. Folate-binding proteins. *Annu. Rev. Nutr.* (1990). doi:10.1146/annurev.nutr.10.1.319 • Abu Seman, N., Wan Mohamad, W. N., Østenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112

GIF

• Ferrand, A. et al. Biochemical and Hematologic Manifestations of Gastric Intrinsic Factor (GIF) Deficiency: A Treatable Cause of B12 Deficiency in the Old Order Mennonite Population of Southwestern Ontario. *JIMD Rep.* (2015). doi:10.1007/8904_2014_351 • Sturm, A. C. et al. Hereditary intrinsic factor deficiency in chaldeans. *JIMD Rep.* (2013). doi:10.1007/8904_2012_133 • Tanner, S. M., Sturm, A. C., Baack, E. C., Liyanarachchi, S. & De La Chapelle, A. Inherited cobalamin malabsorption. Mutations in three genes reveal functional and ethnic patterns. *Orphanet J. Rare Dis.* (2012). doi:10.1186/1750-1172-7-56 • Quadros, E. V. Advances in the understanding of cobalamin assimilation and metabolism. *British Journal of Haematology* (2010). doi:10.1111/j.1365-2141.2009.07937.x • Tanner, S. M. et al. Hereditary juvenile cobalamin deficiency caused by mutations in the intrinsic factor gene. *Proc.Natl.Acad.Sci.U.S.A* (2005). • Agr  us, L. et al. Rationale in diagnosis and screening of atrophic gastritis with stomach-specific plasma biomarkers. *Scand. J. Gastroenterol.* (2012). doi:10.3109/00365521.2011.645501 • Semeco, A. Top 12 Foods That Are High in Vitamin B12. *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>. • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#__text=Vitamin%20B12%20is%20naturally%20found.5%2C13%2D15%5D. • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003). • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011

MTHFD1

• Jiang, J., Zhang, Y., Wei, L., Sun, Z. & Liu, Z. Association between MTHFD1 G1958A polymorphism and neural tube defects susceptibility: A meta-analysis. *PLoS One* (2014). doi:10.1371/journal.pone.0101169 • Field, M. S., Kamynina, E., Watkins, D., Rosenblatt, D. S. & Stover, P. J. Human mutations in methylenetetrahydrofolate dehydrogenase 1 impair nuclear de novo thymidylate biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* (2014). doi:10.1073/pnas.1414555112 • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>. • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998>. • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Abu Seman, N., Wan Mohamad, W. N., Østenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674 • Wu, J. et al. Polymorphisms in MTHFD1 Gene and Susceptibility to Neural Tube Defects: A Case-Control Study in a Chinese Han Population with Relatively Low Folate Levels. *Med. Sci. Monit.* (2015). doi:10.12659/MSM.895155 • Carroll, N. et al. Analysis of the MTHFD1 promoter and risk of neural tube defects. *Hum. Genet.* (2009). doi:10.1007/s00439-008-0616-3 • Neagods, F. C., Cretu, R., Tutulan-Cunita, A., Stoian, V. & Bohiltea, L. C. Methylenetetrahydrofolate dehydrogenase (MTHFD) enzyme polymorphism as a maternal risk factor for trisomy 21: a clinical study. *J. Med. Life* (2010). • Silva, L. M. R. D. da et al. Head and neck carcinogenesis : impact of MTHFD1 G1958A polymorphism. *Head Neck Carcinog.* (2011) • Sutherland, H. G. et al. Association study of MTHFD1 coding polymorphisms R134K and R653Q with migraine susceptibility. *Headache* (2014). doi:10.1111/head.12428 • Saha, T., Dutta, S., Rajamma, U., Sinha, S. & Mukhopadhyay, K. A pilot study on the contribution of folate gene variants in the cognitive function of ADHD probands. *Neurochem. Res.* (2014). doi:10.1007/s11064-014-1393-0 • Zheng, J. et al. MTHFD1 polymorphism as maternal risk for neural tube defects: a meta-analysis. *Neurol. Sci* (2015). • Lorenc, A., Seremak-Mrozkievicz, A., Barlik, M., Wolski, H. & Drews, K. The role of 401a>G polymorphism of methylenetetrahydrofolate dehydrogenase gene (MTHFD1) in fetal hypotrophy. *Ginekol. Pol.* (2014). doi:10.17772/gp/1759 • Cri  an, T. O. et al. The MTHFD1 c.1958 G>A polymorphism and recurrent spontaneous abortions. *J. Matern. Fetal. Neonatal Med.* (2011). doi:10.3109/14767051003702794

MTHFR

• Mischoulon, D. & Raab, M. F. The role of folate in depression and dementia. *Journal of Clinical Psychiatry* (2007). • Goyette, P. et al. Human methylenetetrahydrofolate reductase: Isolation of cDNA, mapping and mutation identification. *Nat. Genet.* (1994). doi:10.1038/ng0694-195 • Hua, Y., Zhao, H., Kong, Y. & Lu, X. Association between Alzheimer's disease and the NOS3 gene Glu298Asp polymorphism. *Int. J. Neurosci.* (2014). doi:10.3109/00207454.2013.834336 • Tran, P. et al. Multiple transcription start sites and alternative splicing in the methylenetetrahydrofolate reductase gene result in two enzyme isoforms. *Mamm. Genome* (2002). doi:10.1007/s00335-002-2167-6 • F  dingner, M., H  rl, W. H. & Sunder-Plassmann, G. Molecular biology of 5,10-methylenetetrahydrofolate reductase. *Journal of Nephrology* (2000). doi:10.5860/CHOICE.39-4838 • E. Trimmer, E. Methylenetetrahydrofolate Reductase: Biochemical Characterization and Medical Significance. *Curr. Pharm. Des.* (2013). doi:10.2174/1381612811319140008 • Wu, X. et al. Association between the MTHFR C677T Polymorphism and Recurrent Pregnancy Loss: A Meta-Analysis. *Genet. Test. Mol. Biomarkers* (2012). doi:10.1089/gtmb.2011.0318 • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Papakostas, G. I. et al. Effect of adjunctive L-methylfolate 15 mg among inadequate responders to SSRIs in depressed patients who were stratified by biomarker levels and genotype: Results from a randomized clinical trial. in *Journal of Clinical Psychiatry* (2014). doi:10.4088/JCP.13m08947 • Schneider, J. A., Rees, D. C., Liu, Y.-T., Clegg, J. B. Worldwide Distribution of a Common Methylenetetrahydrofolate Reductase Mutation. *The American Journal of Human Genetics* 62, 1259–1260 (1998). • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998>. • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • Sibani, S. et al. Characterization of six novel mutations in the methylenetetrahydrofolate reductase (MTHFR) gene in patients with homocystinuria. *Hum. Mutat.* (2000). doi:10.1002/(SICI)1098-1004(200003)15:33.0.CO;2-1 • Goyette, P. et al. Human methylenetetrahydrofolate reductase: Isolation of cDNA, mapping and mutation identification. *Nat. Genet.* (1994). doi:10.1038/ng0694-195 • Abu Seman, N., Wan Mohamad, W. N., Østenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>. • Matthews, R. G. & Daubner, S. C. Modulation of methylenetetrahydrofolate reductase activity by S-adenosylmethionine and by dihydrofolate and its polyglutamate analogues. *Adv. Enzyme Regul.* (1982). doi:10.1016/0065-2571(82)90012-7 • Yamada, K., Strahler, J. R., Andrews, P. C. & Matthews, R. G. Regulation of human methylenetetrahydrofolate reductase by phosphorylation. *Proc. Natl. Acad. Sci.* (2005). doi:10.1073/pnas.0504786102 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Schwahn, B. & Rozen, R. Polymorphisms in the methylenetetrahydrofolate reductase gene: Clinical consequences. *American journal of pharmacogenomics / genomics-related research in drug development and clinical practice* (2001). doi:10.2165/00129785-2001101030-00004 • Yamada, K., Chen, Z., Rozen, R. & Matthews, R. G. Effects of common polymorphisms on the properties of recombinant human methylenetetrahydrofolate reductase. *Proc. Natl. Acad. Sci.* (2001). doi:10.1073/pnas.261469998 • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* (2001). doi:10.1161/01.CIR.103.22.2674 • Bailey, L. B. Folate, Methyl-Related Nutrients, Alcohol, and the MTHFR 677C/T Polymorphism Affect Cancer Risk: Intake Recommendations. *J. Nutr.* (2003). doi:10.1093/jn/133.11.3748S • Nishiyama, M., Kato, Y., Hashimoto, M., Yukawa, S. & Omori, K. Apolipoprotein E, Methylenetetrahydrofolate Reductase(MTHFR) Mutation and the Risk of Senile Dementia. *An Epidemiological Study Using the Polymerase Chain Reaction(PCR) Method.* *J. Epidemiol.* (2000). doi:10.2188/jea.10.163 • Reilly, R., McNulty, H., Pentieva, K., Strain, J. J. & Ward, M. MTHFR 677TT genotype and disease risk: Is there a modulating role for B-vitamins?. *Proc. Nutr. Soc.* (2014). doi:10.1017/S0029665113003613

MTHFS

• Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • Anguera, M. C., & Stover, P. J. (2006). Methenyltetrahydrofolate synthetase is a high-affinity catecholamine-binding protein. *Archives of Biochemistry and Biophysics.* <https://doi.org/10.1016/j.abb.2006.09.008> • Field, M. S., Anderson, D. D., & Stover, P. J. (2011). Mthfs is an essential gene in mice and a component of the purinosome. *Frontiers in Genetics.* <https://doi.org/10.3389/fgene.2011.00036> • Lee, K. M., Lan, Q., Krickler, A., Purdue, M. P., Grulich, A. E., Vajdic, C. M., ... Armstrong, B. K. (2007). One-carbon metabolism gene polymorphisms and risk of non-Hodgkin lymphoma in Australia. *Human Genetics.* <https://doi.org/10.1007/s00439-007-0431-2> • Matakidou, A., El Galta, R., Rudd, M. F., Webb, E. L., Bridle, H., Eisen, T., & Houlston, R. S. (2007). Prognostic significance of folate metabolism polymorphisms for lung cancer. *British Journal of Cancer.* <https://doi.org/10.1038/sj.bjc.6603830> • Rodan, L. H., Qi, W., Ducker, G. S., Demirbas, D., Laine, R., Yang, E., ... Berry, G. T. (2018). 5,10-methylenetetrahydrofolate synthetase deficiency causes a neurometabolic disorder associated with microcephaly, epilepsy, and cerebral hypomyelination. *Molecular Genetics and Metabolism.* <https://doi.org/10.1016/j.ymgme.2018.06.006> • Zhao, R., Chen, Y., Tan, W., Waly, M., Sharma, A., Stover, P. J., ... Deth, R. C. (2001). Relationship between dopamine-stimulated phospholipid methylation and the single-carbon folate pathway. *Journal of Neurochemistry.* <https://doi.org/10.1046/j.1471-4159.2001.00471.x> • Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). *Healthline* (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6.-Citrus-fruits>. • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998>. • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. *Xenobiotica* (2014). doi:10.3109/00498254.2013.845705 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Abu Seman, N., Wan Mohamad, W. N., Østenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. *Clin. Epigenetics* (2015). doi:10.1186/s13148-015-0049-5 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell Metabolism* (2017). doi:10.1016/j.cmet.2016.08.009 • Voutilainen, S.,

MTR

• Semeco, A. Top 12 Foods That Are High in Vitamin B12. Healthline (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>. • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#:~:text=Vitamin%20B12%20is%20naturally%20found,5%2C13%2D15%5D.> • Dutta, S., Shaw, J., Chatterjee, A., Sarkar, K., Usha, R., Chatterjee, A., ... Mukhopadhyay, K. (2011). Importance of gene variants and co-factors of folate metabolic pathway in the etiology of idiopathic intellectual disability. *Nutritional Neuroscience*. <https://doi.org/10.1179/1476830511Y.0000000016> • Ganz, A. B., Shields, K., Fomin, V. G., Lopez, Y. S., Mohani, S., Lovesky, J., ... Caudill, M. A. (2016). Genetic impairments in folate enzymes increase dependence on dietary choline for phosphatidylcholine production at the expense of betaine synthesis. *FASEB Journal*. <https://doi.org/10.1096/fj.201500138RR> • Ono, H., Iwasaki, M., Kuchiba, A., Kasuga, Y., Yokoyama, S., Onuma, H., ... Tsugane, S. (2012). Association of dietary and genetic factors related to one-carbon metabolism with global methylation level of leukocyte DNA. *Cancer Science*. <https://doi.org/10.1111/cas.12013> • Masud, R., & Qureshi, I. Z. (2011). Tetra primer ARMS-PCR relates folate/homocysteine pathway genes and ACE gene polymorphism with coronary artery disease. *Molecular and Cellular Biochemistry*. <https://doi.org/10.1007/s11033-011-1077-7> • Haghir, R., Mashayekhi, F., Bidabadi, E., & Salehi, Z. (2016). Analysis of methionine synthase (rs1805087) gene polymorphism in autism patients in northern Iran. *Acta Neurobiologica Experimentalis*. <https://doi.org/10.21307/ane-2017-030> • Sata, F., Yamada, H., Kishi, R., & Minakami, H. (2012). Maternal folate, alcohol and energy metabolism-related gene polymorphisms and the risk of recurrent pregnancy loss. *Journal of Developmental Origins of Health and Disease*. <https://doi.org/10.1017/S2040174412000359> • Saha, T., Chatterjee, M., Verma, D., Ray, A., Sinha, S., Rajamma, U., & Mukhopadhyay, K. (2018). Genetic variants of the folate metabolic system and mild hyperhomocysteinemia may affect ADHD associated behavioral problems. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. <https://doi.org/10.1016/j.pnpbp.2018.01.016> • Weiner, A. S., Boyarskikh, U. A., Voronina, E. N., Mishukova, O. V., & Filipenko, M. L. (2014). Methylenetetrahydrofolate reductase C677T and methionine synthase A2756G polymorphisms influence on leukocyte genomic DNA methylation level. *Gene*. <https://doi.org/10.1016/j.gene.2013.09.098> • Szczepańska, M., Mostowska, A., Wirstlein, P., Lianeri, M., Marianowski, P., Skrzypczak, J., & Jagodziński, P. P. (2011). Polymorphic variants of folate and choline metabolism genes and the risk of endometriosis-associated infertility. *European Journal of Obstetrics Gynecology and Reproductive Biology*. <https://doi.org/10.1016/j.ejogrb.2011.02.003> • Saha, T., Chatterjee, M., Sinha, S., Rajamma, U., & Mukhopadhyay, K. (2017). Components of the folate metabolic pathway and ADHD core traits: An exploration in eastern Indian probands. *Journal of Human Genetics*. <https://doi.org/10.1038/jhg.2017.23> • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Moore, L. D., Le, T., & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003).

MTRR

• Vaughn, J. D. et al. Methionine synthase reductase 66A->G polymorphism is associated with increased plasma homocysteine concentration when combined with the homozygous methylenetetrahydrofolate reductase 677C->T variant. *J. Nutr.* (2004). doi:134/11/2985 [pii] • Brilakis, E. S., Berger, P. B., Ballman, K. V. & Rozen, R. Methylenetetrahydrofolate reductase (MTHFR) 677C>T and methionine synthase reductase (MTRR) 66A>G polymorphisms: Association with serum homocysteine and angiographic coronary artery disease in the era of flour products fortified with folic acid. *Atherosclerosis* (2003). doi:10.1016/S0021-9150(03)00098-4 • Beyer, K. et al. Methionine synthase polymorphism is a risk factor for Alzheimer disease. *Neuroreport* (2003). doi:10.1097/00001756-200307180-00022 • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#:~:text=Vitamin%20B12%20is%20naturally%20found,5%2C13%2D15%5D.> • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#:~:text=Vitamin%20B12%20is%20naturally%20found,5%2C13%2D15%5D.> • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003). • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Semeco, A. Top 12 Foods That Are High in Vitamin B12. Healthline (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>. • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003). • Moore, L. D., Le, T., & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • MTRR 5-methyltetrahydrofolate-homocysteine methyltransferase reductase [Homo sapiens (human)] - Gene - NCBI. National Center for Biotechnology Information Available at: <https://www.ncbi.nlm.nih.gov/gene/4552>. • Leclerc, D. et al. Cloning and mapping of a cDNA for methionine synthase reductase, a flavoprotein defective in patients with homocystinuria. *Proc. Natl Acad. Sci. USA* (1998). doi:10.1073/pnas.95.6.3059 • Leclerc, D. et al. Cloning and mapping of a cDNA for methionine synthase reductase, a flavoprotein defective in patients with homocystinuria. *Proc. Natl Acad. Sci. USA* (1998). doi:10.1073/pnas.95.6.3059 • Wilson, A. et al. A common variant in methionine synthase reductase combined with low cobalamin (Vitamin B12) increases risk for spina bifida. *Mol. Genet. Metab.* (1999). doi:10.1006/mgme.1999.2879 • James, S. J. et al. Abnormal folate metabolism and mutation in the methylenetetrahydrofolate reductase gene may be maternal risk factors for Down syndrome. *Am. J. Clin. Nutr.* (1999). doi:10.1093/ajcn/70.4.495 • Leclerc, D. et al. Molecular cloning, expression and physical mapping of the human methionine synthase reductase gene. *Gene* (1999). doi:10.1016/S0378-1119(99)00431-X • Doolin, M.-T. et al. Maternal Genetic Effects. Exerted by Genes Involved in Homocysteine Remethylation, Influence the Risk of Spina Bifida. *Am. J. Hum. Genet.* (2002). doi:10.1086/344209 • Pietrzyk, J. J., Bik-Multanowski, M., Sanak, M., & Twardowska, M. Polymorphisms of the 5,10-methylenetetrahydrofolate and the methionine synthase reductase genes as independent risk factors for spina bifida. *J. Appl. Genet.* (2003). • Zhu, H. et al. Homocysteine remethylation enzyme polymorphisms and increased risks for neural tube defects. *Mol. Genet. Metab.* (2003). doi:10.1016/S1096-7192(03)00088-8 • Moore, L. D., Le, T., & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Bosco, P. et al. Methionine synthase (MTR) 2756 (A?G) polymorphism, double heterozygosity methionine synthase 2756 AG/methionine synthase reductase (MTRR) 66 AG, and elevated homocysteinemia are three risk factors for having a child with down syndrome. *Am. J. Med. Genet.* (2003). doi:10.1002/ajmg.a.20234 • Olteanu, H., Wolthers, K. R., Munro, A. W., Scrutton, N. S. & Banerjee, R. Kinetic and Thermodynamic Characterization of the Common Polymorphic Variants of Human Methionine Synthase Reductase†. *Biochemistry* 43, 1988–1997 (2004). • Semeco, A. Top 12 Foods That Are High in Vitamin B12. Healthline (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>.

TCN1

• Semeco, A. Top 12 Foods That Are High in Vitamin B12. Healthline (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>. • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#:~:text=Vitamin%20B12%20is%20naturally%20found,5%2C13%2D15%5D.> • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003). • Moore, L. D., Le, T., & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Remacha, A. F. et al. Role of serum holotranscobalamin (holoTC) in the diagnosis of patients with low serum cobalamin. Comparison with methylmalonic acid and homocysteine. *Ann. Hematol.* (2014). doi:10.1007/s00277-013-1905-z • Irizar, H. et al. Transcriptomic profile reveals gender-specific molecular mechanisms driving multiple sclerosis progression. *PLoS One* (2014). doi:10.1371/journal.pone.0090482 • Johnston, J., Yang-Feng, T. & Berliner, N. Genomic structure and mapping of the chromosomal gene for transcobalamin I (TCN1): Comparison to human intrinsic factor. *Genomics* 12, 459–464 (1992). • Anello, G. et al. Homocysteine and methylenetetrahydrofolate reductase polymorphism in Alzheimer's disease. *Neuroreport* (2004). doi:10.1097/00001756-200404090-00025 • Johnston, J., Bollekens, J., Allen, R. H. & Berliner, N. Structure of the cDNA encoding transcobalamin I, a neutrophil granule protein. *J. Biol. Chem.* (1989). doi:10.1016/j.tetlet.2007.03.117 • Carmel, R., Parker, J. & Kelman, Z. Genomic mutations associated with mild and severe deficiencies of transcobalamin I (haptocorrin) that cause mildly and severely low serum cobalamin levels. *Br. J. Haematol.* (2009). doi:10.1111/j.1365-2141.2009.07855.x • Matteini, A. M. et al. Transcobalamin-II variants, decreased vitamin B12 availability and increased risk of frailty. *J. Nutr. Heal. Aging* (2010). doi:10.1007/s12603-010-0013-1 • Carmel, R. Haptocorrin (transcobalamin I) and cobalamin deficiencies [10]. *Clinical Chemistry* (2007). doi:10.1373/clinchem.2006.078808 • Carmel, R., Parker, J. & Kelman, Z. Genomic mutations associated with mild and severe deficiencies of transcobalamin I (haptocorrin) that cause mildly and severely low serum cobalamin levels. *Br. J. Haematol.* (2009). doi:10.1111/j.1365-2141.2009.07855.x • Bowen, R. A. R. et al. Elevated vitamin B12levels in autoimmune lymphoproliferative syndrome attributable to elevated haptocorrin in lymphocytes. *Clin. Biochem.* (2012). doi:10.1016/j.clinbiochem.2012.01.016 • Furger, E., Frei, D. C., Schibli, R., Fischer, E. & Protá, A. E. Structural basis for universal corrinoid recognition by the cobalamin transport protein haptocorrin. *J. Biol. Chem.* (2013). doi:10.1074/jbc.M113.483271

TCN2

• Regec, A., Quadros, E. V., Platica, O. & Rothenberg, S. P. The cloning and characterization of the human transcobalamin II gene. *Blood* (1995). • Linnebank, M. et al. Association of transcobalamin c. 776C>G with overall survival in patients with primary central nervous system lymphoma. *Br. J. Cancer* (2012). doi:10.1038/bjc.2012.476 • Martinelli, M. et al. A candidate gene study of one-carbon metabolism pathway genes and colorectal cancer risk. *Br. J. Nutr.* (2013). doi:10.1017/S0007114512002796 • Moore, L. D., Le, T., & Fan, G. DNA methylation and its basic function. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.112 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. *Am. J. Hum. Genet.* (2009). doi:10.1016/j.ajhg.2009.02.011 • Koushk, A. et al. Nonsynonymous polymorphisms in genes in the one-carbon metabolism pathway and associations with colorectal cancer. *Cancer Epidemiol. Biomarkers Prev.* (2006). doi:10.1158/1055-9965.EPI-06-0624 • Semeco, A. Top 12 Foods That Are High in Vitamin B12. Healthline (2020). Available at: <https://www.healthline.com/nutrition/vitamin-b12-foods#10.-Fortified-nondairy-milk>. • Mills, J. L. et al. Folate-related genes and omphalocele. *Am. J. Med. Genet.* (2005). doi:10.1002/ajmg.a.30772 • Martinelli, M. et al. Idiopathic pulmonary fibrosis and polymorphisms of the folate pathway genes. *Clin. Biochem.* (2013). doi:10.1016/j.clinbiochem.2012.10.009 • Office of Dietary Supplements - Vitamin B12. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/#:~:text=Vitamin%20B12%20is%20naturally%20found,5%2C13%2D15%5D.> • Miller, A. L. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative Medicine Review* (2003). • Teplitsky, V. et al. Hereditary partial transcobalamin II deficiency with neurologic, mental and hematologic abnormalities in children and adults. *Isr. Med. Assoc. J.* (2003). • Refsum, H., Johnston, C., Guttormsen, A. B. & Nexø, E. Holotranscobalamin and total transcobalamin in human plasma: Determination, determinants, and reference values in healthy adults. *Clin. Chem.* (2006). doi:10.1373/clinchem.2005.054619 • Seetharam, B., Bose, S. & Li, N. Cellular Import of Cobalamin (Vitamin B-12) 1,2. *J. Nutr* (1999). • Winkelmayer, W. C., Skoupy, S., Eberle, C., Födinger, M. & Sunder-Plassmann, G. Effects of TCN2 776C>G on vitamin B, folate, and total homocysteine levels in kidney transplant patients. *Kidney International*. (2004). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15086930> • Hoffbrand, A. V., Tripp, E., Jackson, B. F., Luck, W. E. & Frater-Schroder, M. N. Hereditary Abnormal Transcobalamin II Previously Diagnosed as Congenital Dihydrofolate Reductase Deficiency. *New England Journal of Medicine* 310, 789–790 (1984). • Garg, G. et al. Polymorphisms in transcobalamin II gene is associated with coronary artery disease in Indian population. *Biomarkers* (2012). doi:10.3109/1354750X.2011.642408 • Cascalheira, J. F. et al. Association of the transcobalamin II gene 776C?G polymorphism with Alzheimer's type dementia: dependence on the 5, 10-methylenetetrahydrofolate reductase 1298A?C polymorphism genotype. *Ann. Clin. Biochem.* (2015). doi:10.1177/0004563214561770

MITOCHONDRIA SNP References

ATP5C1

• Yoshida, M., Muneyuki, E. & Hisabori, T. ATP synthase - A marvellous rotary engine of the cell. *Nature Reviews Molecular Cell Biology* (2001). doi:10.1038/35089509 • Gerhard, D. S. et al. The status, quality, and expansion of the NIH full-length cDNA project: The Mammalian Gene Collection (MGC). *Genome Res.* (2004). doi:10.1101/gr.2596504 • Hillier, L. W. et al. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. *Nature* (2005). doi:10.1038/nature03466 • Cross, R. L. Turning the ATP motor. *Nature* (2004). doi:10.1038/427407b • Deloukas, P. et al. The DNA sequence and comparative analysis of human chromosome 10. *Nature* (2004). doi:10.1038/nature02462 • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. *Nature* (2014). doi:10.1038/nature12985 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. *Nature Reviews Immunology* (2011). doi:10.1038/nri2975 • Kann, O. & Kovács, R. Mitochondria and neuronal activity. *American Journal of Physiology - Cell Physiology* (2007). doi:10.1152/ajpcell.00222.2006 • Pizzorno, J. Mitochondria-fundamental to life and health. *Integrative Medicine (Boulder)* (2014). • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria/#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Yan, W. L., Lerner, T. J., Haines, J. L. & Gusella, J. F. Sequence analysis and mapping of a novel human mitochondrial ATP synthase subunit 9 cDNA (ATP5G3). *Genomics* (1994). doi:10.1006/geno.1994.1631

COX5A

• Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014). • Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • Garavito, R. M. & Mulichak, A. M. The Structure of Mammalian Cytochrome c oxidase subunit II gene results in lower levels of the protein and a mitochondrial encephalomyopathy. Am. J. Hum. Genet. (1999). doi:10.1086/302361 • Zee, J. M. & Glerum, D. M. Defects in cytochrome oxidase assembly in humans: lessons from yeast. Biochem. Cell Biol. (2006). doi:10.1139/O06-201 • Kravtsov, Y. et al. Mitochondrial DNA deletions are abundant and cause functional impairment in aged human substantia nigra neurons. Nat. Genet. (2006). doi:10.1038/ng1778 • Jaksch, M. et al. A systematic mutation screen of 10 nuclear and 25 mitochondrial candidate genes in 21 patients with cytochrome c oxidase (COX) deficiency shows tRNA(Ser)(UCN) mutations in a subgroup with syndromal encephalopathy. Journal of Medical Genetics 35, 895–900 (1998).

COX6C

• Lenka, N., Vijayarathy, C., Mullick, J. & Avadhani, N. G. Structural organization and transcription regulation of nuclear genes encoding the mammalian cytochrome c oxidase complex. Prog Nucleic Acid Res Mol Biol (1998). doi:10.1016/S0079-6603(08)60830-2 • Ewing, R. M. et al. Large-scale mapping of human protein-protein interactions by mass spectrometry. Mol. Syst. Biol. (2007). doi:10.1038/msb4100134 • Gerhard, D. S. et al. The status, quality, and expansion of the NIH full-length cDNA project: The Mammalian Gene Collection (MGC). Genome Res. (2004). doi:10.1101/gr.2596504 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Strausberg, R. L. et al. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Proc. Natl. Acad. Sci. U. S. A. (2002). doi:10.1073/pnas.242603899 • Otsuka, M., Mizuno, Y., Yoshida, M., Kagawa, Y. & Ohta, S. Nucleotide sequence of cDNA encoding human cytochrome c oxidase subunit Vic. Nucleic Acids Research 16, 10916–10916 (1988). • Kish, S. J. et al. Decreased brain protein levels of cytochrome oxidase subunits in Alzheimer's disease and in hereditary spinocerebellar ataxia disorders: A nonspecific change? J. Neurochem. (1999). doi:10.1046/j.1471-4159.1999.0720700.x • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014).

NDUFS3

• Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014). • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Benit, P. Mutant NDUFS3 subunit of mitochondrial complex I causes Leigh syndrome. J. Med. Genet. (2004). doi:10.1136/jmg.2003.014316 • Suhane, S., Kanzaki, H., Arumugaswami, V., Murali, R. & Ramanujan, V. K. Mitochondrial NDUFS3 regulates the ROS-mediated onset of metabolic switch in transformed cells. Biol. Open (2013). doi:10.1242/bio.20133244 • Procaccio, V. et al. Human NDUFS3 gene coding for the 30-kDa subunit of mitochondrial complex I: Genomic organization and expression. Mamm. Genome (2000). doi:10.1007/s003350010160

NDUFS7

• Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • Murray, J. et al. The subunit composition of the human NADH dehydrogenase obtained by rapid one-step immunopurification. J. Biol. Chem. (2004). doi:10.1074/jbc.C300064200 • Ugalde, C., Janssen, R. J. R. J., van den Heuvel, L. P., Smetink, J. A. M. & Nijtmans, L. G. J. Differences in assembly or stability of complex I and other mitochondrial OXPHOS complexes in inherited complex I deficiency. Hum. Mol. Genet. (2004). doi:10.1093/hmg/ddh071 • Ota, T. et al. Complete sequencing and characterization of 21,243 full-length human cDNAs. Nat. Genet. (2004). doi:10.1038/ng1285 • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • Smetink, J. & van, L. Human mitochondrial complex I in health and disease. American journal of human genetics. (1999). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10330338>. • Loeffen, J. L. C. M. et al. cDNA of eight nuclear encoded subunits of NADH:ubiquinone oxidoreductase: Human complex I cDNA characterization completed. Biochem. Biophys. Res. Commun. (1998). doi:10.1006/bbrc.1998.9786 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014).

NDUFS8

• West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014). • Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • De Sury, R., Martinez, P., Procaccio, V., Lunardi, J. & Issartel, J. P. Genomic structure of the human NDUFS8 gene coding for the iron-sulfur TYKY subunit of the mitochondrial NADH:ubiquinone oxidoreductase. Gene (1998). doi:10.1016/S0378-1119(98)00275-3 • Su, C. Y., Chang, Y. C., Yang, C. J., Huang, M. S. & Hsiao, M. The opposite prognostic effect of NDUFS1 and NDUFS8 in lung cancer reflects the oncojanus role of mitochondrial complex I. Sci. Rep. (2016). doi:10.1038/srep31357 • Procaccio, V. et al. cDNA sequence and chromosomal localization of the NDUFS8 human gene coding for the 23 kDa subunit of the mitochondrial complex I. Biochimica et Biophysica Acta - Gene Structure and Expression (1997). doi:10.1016/S0167-4781(97)00020-1

SLC19A1

• Link, R. 15 Healthy Foods That Are High in Folate (Folic Acid). Healthline (2020). Available at: <https://www.healthline.com/nutrition/foods-high-in-folate-folic-acid#6..Citrus-fruits.> • Voutilainen, S., Rissanen, T. H., Virtanen, J., Lakka, T. A. & Salonen, J. T. Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. Circulation (2001). doi:10.1161/01.CIR.103.22.2674 • Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. Cell Metabolism (2017). doi:10.1016/j.cmet.2016.08.009 • Abu Seman, N., Wan Mohamud, W. N., Ostenson, C. G., Brismar, K. & Gu, H. F. Increased dna methylation of the slc30a8 gene promoter is associated with type 2 diabetes in a malay population. Clin. Epigenetics (2015). doi:10.1186/s13148-015-0049-5 • Tanaka, T. et al. Genome-wide Association Study of Vitamin B6, Vitamin B12, Folate, and Homocysteine Blood Concentrations. Am. J. Hum. Genet. (2009). doi:10.1016/j.ajhg.2009.02.011 • Moore, L. D., Le, T. & Fan, G. DNA methylation and its basic function. Neuropsychopharmacology (2013). doi:10.1038/npp.2012.112 • Scaglione, F. & Panzavolta, G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. Xenobiotica (2014). doi:10.3109/00498254.2013.845705 • 5-Methyl-tetrahydrofolate. National Center for Biotechnology Information. PubChem Compound Database (2004). Available at: <https://pubchem.ncbi.nlm.nih.gov/compound/135483998>. • Office of Dietary Supplements - Folate. NIH Office of Dietary Supplements (2020). Available at: <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> • Stanisawska-Sachadyn, A. et al. The reduced folate carrier (SLC19A1) c.80G > a polymorphism is associated with red cell folate concentrations among women. Ann. Hum. Genet. (2009). doi:10.1111/j.1469-1809.2009.00529.x • Ananth, C. V. et al. Reduced folate carrier 80A7G polymorphism, plasma folate, and risk of placental abruption. Hum. Genet. (2008). doi:10.1007/s00439-008-0531-7 • Blanton, S. H. et al. Folate pathway and nonsyndromic cleft lip and palate. Birth Defects Res. Part A - Clin. Mol. Teratol. (2011). doi:10.1002/bdra.20740 • Mazumdar, M. et al. Polymorphisms in maternal folate pathway genes interact with arsenic in drinking water to influence risk of myelomeningocele. Birth Defects Res. Part A - Clin. Mol. Teratol. (2015). doi:10.1002/bdra.23399 • James, S. J. et al. A functional polymorphism in the reduced folate carrier gene and DNA hypomethylation in mothers of children with autism. Am. J. Med. Genet. Part B Neuropsychiatr. Genet. (2010). doi:10.1002/ajmg.b.31094 • Clifford, A. J. et al. Single nucleotide polymorphisms in CETP, SLC46A1, SLC19A1, CD36, BCMO1, APOA5, and ABCA1 are significant predictors of plasma HDL in healthy adults. Lipids Health Dis. (2013). doi:10.1186/1476-511X-12-66 • Yee, S. W. et al. SLC19A1 pharmacogenomics summary. Pharmacogenetics and Genomics 20, 708–715 (2010). • Figueiredo, J. C. et al. Genes involved with folate uptake and distribution and their association with colorectal cancer risk. Cancer Causes Control (2010). doi:10.1007/s10552-009-9489-6 • Cho, Y. et al. Association of reduced folate carrier-1 (RFC-1) polymorphisms with ischemic stroke and silent brain infarction. PLoS One (2015). doi:10.1371/journal.pone.0115295 • Matherly, L. H., Wilson, M. R. & Hou, Z. The major facilitative folate transporters solute carrier 19A1 and solute carrier 46A1: Biology and role in antifolate chemotherapy of cancer. Drug Metabolism and Disposition (2014). doi:10.1124/dmd.113.055723 • Coppède, F., Lorenzoni, V. & Migliore, L. The reduced folate carrier (RFC-1) 80A>G polymorphism and maternal risk of having a child with down syndrome: A meta-analysis. Nutrients (2013). doi:10.3390/nu5072551 • O'Byrne, M. R. et al. Association of folate receptor (FOLR1, FOLR2, FOLR3) and reduced folate carrier (SLC19A1) genes with meningioma. Birth Defects Res. A. Clin. Mol. Teratol. (2010). doi:10.1002/bdra.20706 • Coppède, F. et al. A pilot study evaluating the contribution of SLC19A1 (RFC-1) 80G>A polymorphism to Alzheimer's disease in Italian caucasians. Biomed Res. Int. (2014). doi:10.1155/2014/608104

UQCRC2

• Ota, T. et al. Complete sequencing and characterization of 21,243 full-length human cDNAs. Nat. Genet. (2004). doi:10.1038/ng1285 • Strausberg, R. L. et al. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Proceedings of the National Academy of Sciences of the United States of America. (2002). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12477932>. • Hosokawa, Y., Suzuki, H., Toda, H., Nishikimi, M. & Ozawa, T. The primary structure of the precursor to core protein II, a putative member of mitochondrial processing protease family, of rat mitochondrial cytochrome bc1 complex deduced from cDNA sequence analysis. Biochemistry international. (1990). Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2162168>. • Ewing, R. M. et al. Large-scale mapping of human protein-protein interactions by mass spectrometry. Mol. Syst. Biol. (2007). doi:10.1038/msb4100134 • Rual, J. F. et al. Towards a proteome-scale map of the human protein-protein interaction network. Nature (2005). doi:10.1038/nature04209 • Gerhard, D. S. et al. The status, quality, and expansion of the NIH full-length cDNA project: The Mammalian Gene Collection (MGC). Genome Res. (2004). doi:10.1101/gr.2596504 • Kann, O. & Kovács, R. Mitochondria and neuronal activity. American Journal of Physiology - Cell Physiology (2007). doi:10.1152/ajpcell.00222.2006 • Hu, W. H. et al. Identification and characterization of a novel Nogo-interacting mitochondrial protein (NIMP). J. Neurochem. (2002). doi:10.1046/j.1471-4159.2002.00788.x • Hosokawa, Y., Suzuki, H., Toda, H., Nishikimi, M. & Ozawa, T. Complementary DNA encoding core protein II of human mitochondrial cytochrome bc1 complex. Substantial diversity in deduced primary structure from its yeast counterpart. J. Biol. Chem. (1989). • Friedman, J. R. & Nunnari, J. Mitochondrial form and function. Nature (2014). doi:10.1038/nature12985 • West, A. P., Shadel, G. S. & Ghosh, S. Mitochondria in innate immune responses. Nature Reviews Immunology (2011). doi:10.1038/nri2975 • MitoQ. Which Foods Help Your Mitochondria? MitoQ (2019). Available at: <https://www.mitoq.com/blog/which-foods-help-your-mitochondria#:~:text=Simple carbohydrates like white flour,efficient, long-lasting way.> • Pizzorno, J. Mitochondria-fundamental to life and health. Integrative Medicine (Boulder) (2014).